



VA: VGH / UBCH / GFS
VC: BP / Purdy / GPC

ORDERS

ADDRESSOGRAPH

COMPLETE OR REVIEW ALLERGY STATUS PRIOR TO WRITING ORDERS

EYE CARE CENTRE SURGERY ORDERS

(items with check boxes must be selected to be ordered)

(Page 1 of 1)

Date: _____ Time: _____

Level of Anxiety: _____ Low _____ Moderate _____ High

MONITORING:

- Vital signs pre-op and at end of surgery: BP, HR, RR, sedation scale, O₂ saturation
- Five minutes post administration of benzodiazepine or narcotic: BP, HR, RR, sedation scale, O₂ saturations Q15MIN

PATIENT PREPARATION:

Saline lock PRN

Operative eye(s): right left

For intraocular procedures administer 1 drop of the following preoperative eye drops to operative eye Q15MIN x 3 upon admission:

- tetracaine 0.5%
- cyclopentolate 1%
- ofloxacin 0.3%
- diclofenac 0.1% (VOLTAREN)
- PHENYLEphrine 2.5% *OR* PHENYLEphrine 10%
- tropicamide 1%
- pilocarpine 2%

Intraocular procedures:

- Prepare lashes with povidone-iodine 10% solution
- Patients with allergy to seafood or iodine: use chlorhexidine 0.05% (1:2000) solution

Plastic procedures:

Use chlorhexidine 0.05% (1:2000) solution as per protocol

Anxiety management:

- lorazepam 0.5 to 1 mg SL PRN to a maximum of 1 mg total cumulative dose *OR*
- midazolam protocol:
midazolam 0.5 to 1 mg IV initially, followed by
midazolam 0.5 to 1 mg IV Q2MIN to a maximum cumulative dose of 3 mg

Pain management:

- fentanyl protocol:
fentanyl 25 mcg IV initially, followed by
fentanyl 25 mcg IV Q2MIN to a maximum cumulative dose of 100 mcg
- acetaminophen 325 to 650 mg Q4H PO PRN

DISCHARGE:

Patient may be discharged when sedation and discharge criteria are met and at least 30 minutes post administration of any benzodiazepine or narcotic.

Time
Processed
RN/LPN
Initials
Comments

Prescriber's Signature
ECCSP

Printed Name
VCH.VA.PPO.260 | Rev.JUN.2014

College ID